

GLOBAL JOURNAL OF ENGINEERING SCIENCE AND RESEARCHES
RECENT ADVANCES IN SYNTHESIS OF 2H-INDAZOLE FROM 2-SUBSTITUTED
BENZALDEHYDE: A REVIEW


Pravinsing S. Girase¹, Bhikan J. Khairnar¹ & Bhata R. Chaudhari^{1,2}

¹Department of Chemistry, JET's Z. B. Patil College, Dhule (MS), India.

²Department. of Chemistry, SSVPS's ACS College, Shindkheda, Dhule (MS), India

ABSTRACT

Several compounds has more than one nitrogen contained bicyclic heterocycles were important structural motifs found in numerous natural products and bioactive molecules. In recent years, the synthesis of bicyclic compounds possessing imidazole heterocycle as central core has been the focus of great interest. They are associated with diversified biological activities.



I. INTRODUCTION

Heterocycles are broadly found in naturally occurring compounds and have an immense importance in the design and discovery of new compounds for pharmaceutical applications [1]. Among the numerous heterocycles, nitrogen containing heterocyclic skeleton show a key role in alkaloids chemistry. These alkaloid compounds are gaining a lot of importance and they have exhibit a broad spectrum of pharmacological and biological activities [2]. Over the past decades, a great deal of research interests has been devoted toward the synthesis of heterocyclic compounds because of their significant biological activities [3]. Indazole is a ten- π electron aromatic heterocyclic system in which pyrazole ring condensed with the benzene ring [4]. Indazole also called benzpyrazole or isoindazole is a heterocyclic aromatic organic compound. Comparable the pyrazole molecule indazole resembles both pyridine and pyrrole and its reactivity reproduces this twofold activity [5]. The indazole ring has two nitrogen atoms and presents annular tautomerism with regards to the position of the NH hydrogen atom [6]. There are two tautomeric forms for indazole, the 1*H*- and 2*H*- form (Figure 1).




Figure 1: Tautomers of Indazole.

he synthesis of bicyclic pyrazole bearing heterocyclic compounds 2*H*-Indazole unit has been reorganized by a "privileged structure" and is an important pharmacophore in medicinal chemistry [7]. Indazoles are rare in nature and to date only three natural products possessing the indazole ring have been isolated are Nigellidine, Nigeglanine and Nigellidine (Figure 2) [8].




Figure 2: Natural products possessing the indazole ring




Figure 3: The structure of approved drugs having indazole cores and their corresponding pharmaceutical properties. Interestingly, in the field of drug discovery, the Indazole scaffolds received a lot of interest because they act as efficient bioisosters of indole and benzimidazole [7, 9]. Molecules constitute 2H-Indazole skeleton are found the key subunit in various drug substances has been receiving significant attention due to its pronounced biological activities, including anti-inflammatory [10], anti-platelet [11], antidepressant [12], anticancer [13], anti-tumor [14], anti-HIV [15], antimicrobial [16] and anti-spermatogenic [17] activities. Some 2H-indazole derivatives are proved to have high affinity for the imidazoline I₂ receptor [18], potent affinity for estrogen β -receptor [19] and 5-HT_{1A} receptors [20]. Some clinically approved important drugs such as lonidamine [21], bendacort [22], bendazac [23], granisetron [24], axitinib [25], pazopanib [26], niraparib (MK-4827) [27], YC-1 [28], selective estrogen β -receptor (Figure 3 (I)) [19] and farnesoid-X receptor agonist (Figure 3 (II)) [29] were known to involve the indazole and its

related compounds have been manufactured and used worldwide (**Figure 3**). The indazole ring system is also present in many other compounds such as herbicides [30], dyes-pigments [31] and as optoelectronic chromophores used in fabrication of devices like OLEDs [32].

II. REVIEW OF LITERATURE


The tautomeric equilibrium between *1H*- and *2H*-indazole has been calculated by thermo-chemical and photo physical techniques. The results shows *1H* tautomer more stable than *2H* tautomer, for example 1-methyl-*1H*-indazole (3.2 kcal/mol) [33] more stable and stronger base [34] than 2-methyl-*2H*-indazole. Dipole moment of *2H* tautomer is more than *1H* tautomer [35]. (**Table 1**)

Table 1: Basicity and dipole moment comparison of indazole tautomers

Physical technique	1-methyl- <i>1H</i> -indazole	2-methyl- <i>2H</i> -indazole
Basicity (P^{kb})	0.42	2.02
Dipole moment (D)	1.50	3.4

The chemistry of *2H*-indazoles has not been explored as well as the chemistry of *1H*-indazoles. However, the discovery that *N*-2 substituted *2H*-indazole compounds exhibit biological activities, which has produced recent interest in their simple and efficient preparation. Although a number of methods were known for the synthesis of Indazoles, amongst them most of the existing approaches are targeted *1H*-Indazole or mixture of both *1H*- and *2H*-Indazoles. *2H*-Indazoles are less studied as compared to *1H*-Indazole; due to the difficulty in their preparation. Therefore the formation of *2H*-Indazoles still remains a challenging task [7].

In past, few synthetic approaches have been developed for the synthesis of *2H*-indazoles (**Scheme 1**), which can be achieved by the different synthetic methodologies viz. C-C and C-N bond formation using different catalysts through the one pot three-component condensation reaction of 2-bromobenzaldehyde, primary amine and salt of azide (**route a**) [36-42]. A copper catalyst plays the key role in both C-N bond formation between the aryl bromide and the azide and N-N bond formation between the imine and the azide. The condensation reaction of 2-azidobenzaldehyde with primary amine via formation of imine as intermediate in situ (**route b**) [43, 44]. One traditional approach to yield *2H*-indazole is intramolecular reductive *N*-hetero-cyclization of *N*-(2-nitrobenzylidene)anilines formed from 2-nitro benzaldehyde with primary amine (**route c**) [45-51].




Scheme 1: Literature available methods for the synthesis of 2H-Indazoles.

In other way arynes with easily obtained sydnone by a sequence involving [3+2] dipolar cycloaddition reaction in presence of TBAF (1.6 equiv) (**route d**) [52]. Other methods involve the reaction of bis(2-chloromethyl)arylzinc reagents with aryldiazonium salt in THF:NMP (4:1) (**route e**) [53]. Song and Yee reports that the Pd(OAc)₂/dppf as homogeneous catalytic system and NaO^tBu as base through intramolecular amination reaction of *N*-aryl-*N*-(*o*-bromobenzyl)-hydrazines (**route f**) [54]. Kim and his research group reported the two step procedure for the synthesis of 2*H*-indazole via cross coupling reaction of aryl hydrazine with the Baylis-Hillman adduct (**route g**) [55]. In another reported process reactions of aryl hydrazine with 2-halophenyl acetylenes (**route h**) [56]. Some reports are shows that palladium-catalyzed intramolecular C-N bond formation of *o*-alkyne azoarene (**route i**) [57] and iodine mediated intramolecular oxidative annulation of benzyl sp³ C-H functionalization of *o*-alkyl azoarenes (**route j**) [58]. Also Rh (III)-catalyzed direct addition of an azobenzene C-H bond to an aldehyde with subsequent cyclization and aromatization (**route k**) [59]. A Rh(III)-catalyzed tandem C-H alkylation as well as intramolecular decarboxylative cyclization of azoxy compounds with diazoesters (**route l**) [60] or alkynes (**route m**) [61] via a classic [4 + 1] annulation manner. The 2*H*-indazoles are synthesis by Davis-Beirut reaction from *o*-nitroso benzaldehyde and primary amines under both acid and base catalysis [62]. Base-catalysed reaction of 2-nitrobenzyl triphenylphosphonium bromide with Aryl isocyanate [63] and Fe-catalysed N-N bond formation of 2-azidophenyl ketoximes [64].

Some selected reactions for the synthesis of 2*H*-indazoles are described here.

Emil Fischer and Kuzel (1880) have reported the first synthesis of heterocycle containing an indazole ring by heating *o*-hydrazinocinnamic acid [7]. In an attempt to prepare the anhydride of *o*-hydrazinocinnamic acid, Fisher

got many products, out of these products one of which was a new substance containing no oxygen. The analysis of the reaction mixture demonstrated the elimination of an acetic acid molecule and led to the conclusion that the structure of the new substance was a pyrazole ring condensed with a benzene ring which requires longer reaction time and low to moderate yield of product. Fischer and Kuzel described this transformation as “remarkable of the highest degree” and named the product “indazole” by analogy with the previously known indole (Scheme 2).




Scheme 2: First reported synthesis of Indazole

Synthesis of 2H-indazole from 2-substituted benzaldehyde (Route a-c):

1) M. R. Kumar *et al.* (2011) [36]


M. R. Kumar *et al.* have reported the synthesis of 2-aryl-2H-Indazoles through one-pot, three-component condensation reactions of 2-bromobenzaldehydes, primary amines, and sodium azide by using CuI (10 mol%)/TMEDA (10 mol%) as a homogeneous catalyst system in DMSO at 120°C for 12h resulted in good yields (Scheme 3).



Scheme 3

2) B. M. Reddy and co-workers (2013) [37]

B. M. Reddy and co-workers described that a synthesis of 2H-indazoles was achieved from 2-bromobenzaldehydes, primary amines and sodium azide through consecutive condensation, C–N and N–N bond formations catalyzed by a heterogeneous Cu(II)–HT (20mg, Cu:Al ratio 3:1) catalyst in DMSO at 120°C temperature for 6 h (Scheme 4). A series of 2H-indazoles were synthesised in good to excellent yields. Furthermore, the Cu(II)–HT catalyst can be reused for three runs without any significant loss of activity.




Scheme 4

3) N. Khatunetal. (2014) [38]

N. Khatunetal. have reported a CuO (2.5 mol%) nano catalysed one-pot synthesis of 2H-indazoles has been devised from 2-bromobenzaldehydes, primary amines and sodium azide with Cs₂CO₃ (2 equiv.) as base in DMSO at 120°C


under ligand free conditions. This catalytic system gave the yield in the range of 40-84%. The catalyst can be recycled up to three times (Scheme 5).



Scheme 5

4) R. K. Sodhietal. (2014) [39]


R. K. Sodhietal. have reported the one pot multicomponent synthesis of 2H-indazole from 2-bromobenzaldehyde, primary amines and sodium azide through consecutive C-N and N-N bond forming reaction was carried out in the presence of different amine functionalized silica/starch-M(acac)_n catalysts. ASS-Cu(acac)₂ (0.2 g, 1.1 wt% Cu) was turned out to be the best catalyst in DMSO at 100°C for the synthesis of 2H-indazoles (Scheme 6). It was found that ASS-Cu(acac)₂ could be recycled for four times.



Scheme 6

5) H. Sharghietal. (2017) [40]


H. Sharghietal. have reported the synthesis of 2H-indazole via one pot multicomponent condensation of 2-bromobenzaldehyde, primary amine with sodium azide by using salen. The support-free Cu(II)-salen complex (7 mol%) as a catalyst under nearly solvent-free conditions at 120°C (Scheme 7).



Scheme 7

6) M. N. Soltani Rad (2017) [41]


M. N. Soltani Rad has reported an ultrasonic promoted facile and convenient one-pot three-component reaction of 2-bromobenzaldehyde, primary amine and tetra-butyl ammonium azide (TBAA) used as an azide source in presence of copper-doped silica cuprous sulphate (CDSCS) in DMSO at room temperature furnishes 2H-indazoles in good yields (Scheme 8).



Scheme 8

7) S. Behrouz (2016) [42]


S. Behrouz reported the multicomponent synthesis of 2H-indazole from condensation of 2-bromobenzaldehyde, structurally diverse amines, and [bmim]N₃ is used as an azide source in the presence of Cu/aminoclay/reduced graphene oxide nanohybrid (Cu/AC/r-GO nanohybrid) as heterogeneous catalyst in DMF at 110°C gave 2H-indazoles in good yields (Scheme 9).



Scheme 9

8) D. S. Sharada and group (2014-2017) [43]


D. S. Sharada and group have described a catalyst-free, solvent-free (neat condition) and microwave-irradiated approach at 110°C for 50-60 min, to 2H-indazoles through consecutive formation of C=N and N=N bonds from 2-azidobenzaldehyde with various primary amines like aromatic, heteroaromatic, aliphatic and benzylic amines which offered wide range of 2H-indazoles in excellent yields (Scheme 10).



Scheme 10

9) J. Hu *et al.* (2011) [44]


J. Hu *et al.* have developed an intramolecular amination reaction of 2-azidobenzaldehyde with primary amine under mild condition. By employing a combination of CuI (0.1 equiv.) with tertiary amines TMEDA or TEA (0.5 equiv.) which is homogeneous catalytic system in THF at rt for 4h or at 40°C for 2h used for the intramolecular N–N bond formation reaction (Scheme 11).



Scheme 11

10) Akazomeetal. (1994) [45]


Akazomeetal. demonstrated the palladium-catalyzed intramolecular reductive *N*-heterocyclization of (2-nitrobenzylidene) amines to give the corresponding 2*H*-indazoles. Reactions were conducted in a stainless reactor under 20 kg cm⁻² of initial CO pressure in the presence of a PdCl₂(PPh₃)₂ (5 mol%) and SnCl₂ (50 mol%) at 100°C for 16h (Scheme 12). Carbon monoxide effectively worked as a deoxygenating agent of the nitro group which gave a nitrene intermediate. Also, it required high catalyst loading with no recyclability.



Scheme 12

11) J.I.G. Cadogan and R. K. Mackie (1968) [46]


J.I.G. Cadogan and R.K. Mackie reported the new catalytic method for the synthesis of 2-aryl-2*H* indazole derivatives through reductive cyclocondensation of 2-nitrobenzaldehyde with aromatic primary amines in presence of P(OEt)₃ as catalyst at 150°C temperature by Cadogan reaction (Scheme 13).



Scheme 13

12) N. E. Genungetal. (2014) [47]


N. E. Genungetal. reported the synthesis of 2*H*-indazoles by one-pot condensation Cadogan reductive cyclization. As condensation of aryl aldehyde and aniline occurs readily in *iso*-PrOH at elevated temperature, it was envisioned that simply adding tri-*n*-butylphosphine after the condensation was completed, followed by heating, would afford the desired product without the complicating factor of isolating an *o*-iminonitrobenzene intermediate. Fortunately, this method proved effective as the one-pot process afforded indazole in 85% yield (Scheme 14).



Scheme 14

13) T. V. Nykazaetal. (2017) [48]


T. V. Nykazaetal. have reported a readily accessible phosphetane which was a suitable catalyst for the Cadogan indazole synthesis. A small-ring phosphacycle, 1,2,2,3,4,4-hexamethyl-phosphetane, was found to catalyze deoxygenative N-N bond-forming Cadogan heterocyclization of o-nitrobenzaldehydes, o-nitroazobenzenes and related substrates in the presence of hydrosilane terminal reductant. The method provides a simple phosphacatalytic approach to a valuable N-N bond-forming mode (Scheme 15).



Scheme 15

14) A. K. Saikiaetal. (2014) [49]


A. K. Saikiaetal. have reported the effective synthesis of 3-(Arylethynyl)-2H-indazoles through one pot three component condensation using 2-nitroarylaldehydes, primary amines and alkynes co-catalyzed by copper(I) bromide and zinc(II) triflate in toluene refluxed for 24 to 48h (Scheme 16).



Scheme 16

15) N. Sudhapriyaetal. (2014 & 2017) [50]

N. Sudhapriyaetal. reported the synthesis of 3-Alkynyl-2-Aryl-2H-Indazole or 2-aryl-2H-indazole-3-phosphonates via SnCl₂.2H₂O mediated A³-coupling followed by reductive cyclization of 2-nitrobenzaldehyde, primary amine and phenyl acetylene or dialkyl phosphonates (Scheme 17). A rapid and efficient synthesis of indazole through N-N bond formation. This protocol proceeds wide substrate scope and good yields.



Scheme 17

III. CONCLUSION

In this review, we have reporting various methods for the synthesis of 2H-indazoleheterocyclic scaffoldthrough C-N bond formation reactions.

IV. ACKNOWLEDGEMENTS

The author BJK is greatly thankful to Council of Scientific and Industrial Research (CSIR), NewDelhi, India for providing the research fellowship. The authors are also thankful to the Principal, JET's Z.B. Patil College, Dhule for providing the laboratory facilities for this work.

REFERENCES

1. a) A. R. Katritzky, C. W. Rees, *Comprehensive Heterocyclic Chemistry*; Eds. Pergamon Press: New York, 1984, 1. (b) A. R. Katritzky, C. W. Rees, E. F. V. Scriven, *Comprehensive Heterocyclic Chemistry II*; Eds. Pergamon: New York, 1996, 1. (c) F. Couty, G. Evano, *Comprehensive Heterocyclic Chemistry*, Elsevier, Oxford, 2008. (d) E. G. Brown, *Ring nitrogen and key biomolecules: The biochemistry of N-heterocycles*, Kluwer academic, Boston, 1998.
2. G. A. Cordell, *The Alkaloids; Chemistry and Biology*, Volume 69, Ed., 2010, Elsevier, and previous volumes in the series.
3. (a) W. Stadlbauer, *Science of Synthesis*; Georg Thieme: Stuttgart, 2002, 12, 227. (b) M. J. Haddadin, W. E. Conrad, M. J. Kurth, *The Davis-Beirut Reaction: a novel entry into 2H-indazoles and indazolones. Recent biological activity of indazoles.*, *Mini Rev Med Chem*, 2012, 12, 1293.
4. A. A. Hassan, A. A. Aly, H. N. Tawfeek, *Chapter Three - Indazoles: Synthesis and Bond-Forming Heterocyclization*, *Adv. Heterocycl. Chem.* 2018, 125, 235.
5. D. J. Triggle, J. B. Taylor, *Comprehensive Medicinal Chemistry-II*, Ed., Elsevier, 2006.
6. A. T. Balaban, D. C. Oniciu, A. R. Katritzky, *Aromaticity as a Cornerstone of Heterocyclic Chemistry*, *Chem. Rev.* 2004, 104, 2777.
7. (a) A Schmidt, A. beutler, B. Snovydovych, *Recent Advances in the Chemistry of Indazoles*, *Eur. J. Org. Chem.* 2008, 4073. (b) N. Cankařová, J. Hlaváč, V. Krchňák, *Handbook of Reagents for Organic Synthesis*, *Org. Prep. Proced. Int.* 2010, 42, 433. (b) D. D. Gaikwad, A. D. Chapolikar, C. G. Devkate, K. D. Warad, A. P. Tayade, R. P. Pawar, A. J. Domb. *Synthesis of indazole motifs and their medicinal importance: an overview*. *Eur. J. Med. Chem.* 2015, 90, 707.
8. Y. M. Liu, J. S. Yang, Q. U. Liu, *A New Alkaloid and Its Artificial Derivative with an Indazole Ring from Nigella glandulifer*, *Chem. Pharm. Bull.*, a, 2004, 52, 454.
9. L. A. Clutterbuck, C. G. Posada, C. Visintin, D. R. Riddal, B. Lancaster, P. J. Gane, J. Garthwaite, D. L. Selwood, *Oxadiazolyindazole Sodium Channel Modulators are Neuroprotective toward Hippocampal Neurons*, *J. Med. Chem.* 2009, 52, 2694.
10. (a) G. Picciola, F. Ravenna, G. Carenini, P. Gentili, M. Riva, *Heterocyclic compounds containing the residue of a 4-aminophenylalkanoic acid with potential anti-inflammatory activity. IV. Derivatives of 2-phenyl-2H-indazole*, *Farmaco, Ed. Sci.* 1981, 36, 1037. (b) G. A. Bistochi, G. De Meo, M. Pedini, A.

- Ricci, H. Brouilhet, S. Bucherie, M. Rabaud, P. Jacquignon, *Farmaco. N1-substituted 1H-indazole-3-ethyl carboxylates and 1H-indazole-3-hydroxamic acids*, *Ed. Sci.*1981, 36, 315.
11. F.-Y. Lee, J.-C. Lien, L.-J. Huang, T.-M. Huang, S.-C. Tsai, C.-M. Teng, C.-C. Wu, F.-C. Cheng, S.-C. Kuo, *Design, Synthesis, and Preclinical Characterization of Novel, Highly Selective Indole Estrogens*, *J. Med. Chem.*2001, 44, 3747.
 12. Y. Ykeda, N. Takano, H. Matsushita, Y. Shiraki, T. Koide, R. Nagashima, Y. Fuzimura, M. Shindo, S. Suzuki, T. Iwasaki, *Pharmacological studies on a new thymoleptic antidepressant, 1-[3-(dimethylamino)propyl]-5-methyl-3-phenyl-1H-indazole (FS-32)*, *Arzneim. Forsch.*1979, 29, 511.
 13. M. De Lena, V. Lorusso, A. Latorre, G. Fanizza, G. Gargano, L. Caporusso, M. Guida, A. Catino, E. Crucitta, D. Sambiasi, A. Mazzei, *Paclitaxel, cisplatin and lonidamine in advanced ovarian cancer. A phase II study.*, *Eur. J Cancer*2001, 37, 364.
 14. P. G. Baraldi, G. Balboni, M. G. Pavani, G. Spalluto, M. A. Tabrizi, E. De Clercq, J. Balzarini, T. Bando, H. Sugiyama, R. Romagnoli, *Design, Synthesis, DNA Binding, and Biological Evaluation of Water-Soluble Hybrid Molecules Containing Two Pyrazole Analogues of the Alkylating Cyclopropylpyrroloindole (CPI) Subunit of the Antitumor Agent CC-1065 and Polypyrrole Minor Groove Binders*, *J. Med. Chem.*2001, 44, 2536.
 15. W. Han, J. C. Pelletier, C. N. Hodge, *Tricyclic ureas: a new class of HIV-1 protease inhibitors.*, *Bioorg. Med. Chem. Lett.*1998, 8, 3615.
 16. X. Li, S. Chu, V. A. Feher, M. Khalili, Z. Nie, S. Margosiak, V. Nikulin, J. Levin, K. G. Parankle, M. E. Tedder, R. Almassy, K. Appelt, K. M. Yager, *Structure-Based Design, Synthesis, and Antimicrobial Activity of Indazole-Derived SAH/MTA Nucleosidase Inhibitors*, *J. Med. Chem.*2003, 46, 5663.
 17. G. Corsi, G. Palazzo, C. Germani, P. S. Barcellona, B. Silvestrini, *1-Halobenzyl-1H-indazole-3-carboxylic acids. A new class of antispermatogenic agents*, *J. Med. Chem.*1976, 19, 778.
 18. F. Saczewski, J. Saczewski, A. L. Hudson, R. J. Tyacke, D. J. Nutt, J. Man, P. Tabin, *2-(4,5-dihydro-1H-imidazol-2-yl)indazole (indazim) derivatives as selective I(2) imidazoline receptor ligands.*, *Eur J Pharm Sci*, 2003, 20, 201.
 19. M. D. Angelis, F. Stossi, K. A. Carlson, B. S. Katzenellenbogen, J. A. Katzenellenbogen, *Indazole Estrogens: Highly Selective Ligands for the Estrogen Receptor β* , *J. Med. Chem.*, 2005, 48,1132.
 20. S. Andronati, V. Sava, S. Makan, G. Kolodeev, *Synthesis of 3-aryl-1-[(4-phenyl-1-piperazinyl)butyl]indazole derivatives and their affinity to 5-HT1A serotonin and dopamine D1 receptors*, *Pharmazie*, 1999, 54, 99.
 21. (a) S. Di Cosimo, G. Ferretti, P. Papaldo, P. Carlini, A. Fabi, F. Cognetti, *Lonidamine: efficacy and safety in clinical trials for the treatment of solid tumors*. *Drugs Today(Barc)*.2003, 39, 157. (b) A. Caputo, B. Silvestrini, *Lonidamine, a new approach to cancer therapy*, *Oncology* 1984, 41, 2.(c) A. P. Rajput, D. V. Nagarale, *Synthesis, Characterization and Antimicrobial study of piperidine-2,6-diones derivatives*, *Der Pharma Chemica*, 2016, 8(8):182-186.
 22. a)L. Baiocchi. *US Patent 4,001,219; January 4, 1977.*b) (d) A. P. Rajput, D. V. Nagarale, *Synthesis Of 2,2'-{[2,6-Dichloro-1-(Substitutedphenyl)-1,4-Dihydropyridine-3,5-Diyl] Dimethyllydene} Dipropanedinitrile Using Green Catalyst L-Proline-Fe₃O₄ MNP*, *Journal of Applicable Chemistry*, 2018, 7 (6): 1592-1596.
 23. a)H. Cerecetto, A. Gerpe, M. Gonzalez, V. J. Aran, C. O. de Ocariz, *Pharmacological properties of indazole derivatives: recent developments.*, *Mini. Rev. Med. Chem.* 2005, 5, 869. b) A. P. Rajput, D. V. Nagarale, *Thermal Condensation of Enaminoimine Hydrochlorides of 2,6-Dichloro-3,5-diformyl (N-Substituted Phenyl)pyridines*, *Chem Sci Trans.*, 2016, 5(4), 912-917.
 24. P. J. Hesketh, D. R. Gandara, *A possible role for cisplatin in the transient hepatic enzyme* *J. Natl. Cancer Inst.*1991, 83(45), 613.
 25. N. Ferrara, *Vascular endothelial growth factor as a target for anticancer therapy* *Oncologist*, 2004, 9(suppl. 1), 2.
 26. Y. Jia, J. Zhang, J. Feng, F. Xu, H. Pan and W. Xu, *Design, synthesis and biological evaluation of pazopanib derivatives as antitumor agents*, *Chem. Biol. Drug Des.* 2014, 83, 306.
 27. P. Jones, S. Altamura, J. Boueres, F. Ferrigno, M. Fonsi, C. Giomini, S. Lamartina, E. Monteagudo, J. M. Ontoria, M. V. Orsale, M. C. Palumbi, S. Pesci, G. Roscilli, R. Scarpelli, C. Schultz-

- Fademrecht, C. Toniatti, M. Rowley, Discovery of 2-{4-[(3S)-Piperidin-3-yl]phenyl}-2H-indazole-7-carboxamide (MK-4827): A Novel Oral Poly(ADP-ribose)polymerase (PARP) Inhibitor Efficacious in BRCA-1 and -2 Mutant Tumors, *J. Med. Chem.* 2009, 52, 7170.
28. A. Takeuchi, M. Hori, S. Sato, H. S. Ban, T. Kuchimaru, S. Kizaka-Kondoh, T. Yamori, H. Nakamura, Synthesis and biological activity of furanylindazoles as inhibitors of hypoxia inducible factor (HIF)-1 transcriptional activity, *Med. Chem. Commun.* 2012, 3, 1455.
 29. G. M. Benson, K. Bleicher, S. Feng, U. Grether, B. Kuhn, R. E. Martin, J.-M. Plancher, H. Richter, M. Rudolph, S. Taylor, US 2010/0076026.
 30. A. D. Mills, P. Maloney, E. Hassanein, M. J. Haddadin, M. J. Kurth, Synthesis of a Library of 2-Alkyl-3-alkyloxy-2H-indazole-6-carboxamides, *J. Comb. Chem.* 2007, 9, 171.
 31. Ullmann's Encyclopedia of Industrial Chemistry, 7th ed. Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim, 2005.
 32. Y. Zhang, X. Ma, Y. Chen, X. Chen, L. Guo, W. Cao, J. Chen, M.S. Wong, Synthesis and Characterization of Oxadisilole-Fused 1H-Benzo[f]indazoles and 1H-Naphtho[2,3-f]indazoles, *Eur. J. Org. Chem.* 2013, 3005.
 33. J. Catalán, J. C. del Valle, R. M. Claramunt, G. Boyer, J. Laynez, J. Gómez, P. Jiménez, F. Tomás, J. Elguero, Acidity and Basicity of Indazole and its N-Methyl Derivatives in the Ground and in the Excited State, *J. Phys. Chem.* 1994, 98, 10606.
 34. (a) J. Elguero, A. Fruchier, R. Jacquier, *Bull. Soc. Chem. Fr.* 1967, 2619. (b) G. A. Jaffari, A. J. Nunn, *J. Chem. Soc. Perkin Trans*, 1973, 1, 2371. (c) J. Catalan, J. Elguero, *J. Heterocycl. Chem.* 1984, 21, 269. (d) A. P. Rajput, D. V. Nagarale, *Modern Synthetic Tool L-Proline as an Organocatalyst, Journal of Chemical and Pharmaceutical Research*, 2016, 8(7):557-575.
 35. (a) A. Escande, J. Lapasset, R. Faure, E. Vincent, J. Elguero, *Tetrahedron*, 1974, 30, 2903. (b) C. Ogretir, N. Funda (Kaypak) Tay, *J. Mol. Struct: Theochem*, 2002, 588, 145. (c) P. Maunet, J. P. Fayet, M. Fabre, *Bull. Soc. Chem. Fr.* 1975, 1675.
 36. M. R. Kumar, A. Park, N. Park, S. Lee, Consecutive Condensation, C–N and N–N Bond Formations: A Copper- Catalyzed One-Pot Three-Component Synthesis of 2H-Indazole, *Org. Lett.* 2011, 13, 3542.
 37. A. N. Prasad, R. Srinivas, B. M. Reddy, CuII-hydrotalcite catalyzed one-pot three component synthesis of 2H-indazoles by consecutive condensation, C–N and N–N bond formations, *Catal. Sci. Technol.* 2013, 3, 654.
 38. N. Khatun, A. Gogoi, P. Basu, P. Das, B. K. Patel, CuO nanoparticle catalysed synthesis of 2H-indazoles under ligand free conditions, *RSC Adv.*, 2014, 4, 4080.
 39. R. K. Sodhi, A. Changotra, Metal Acetylacetonates Covalently Anchored onto Amine Functionalized Silica/Starch Composite for the One-Pot Thioetherification and Synthesis of 2H-Indazoles, *Catal. Lett.*, 2014, 144, 1819.
 40. H. Sharghi, M. Aberi, P. Shiri, Highly reusable support-free copper(II) complex of para-hydroxy-substituted salen: Novel, efficient and versatile catalyst for C–N bond forming reactions, *Appl. Organometal. Chem.* 2017, e3761.
 41. M. N. Soltani Rad. Ultrasound promoted mild and facile one-pot, three component synthesis of 2H-indazoles by consecutive condensation, CN and NN bond formations catalysed by copper-doped silica cuprous sulphate (CDSCS) as an efficient heterogeneous nano-catalyst., *Ultrason. Sonochem.* 2017, 34, 865.
 42. S. Behrouz, Ultrasound promoted mild and facile one-pot, three component synthesis of 2H-indazoles by consecutive condensation, CN and NN bond formations catalysed by copper-doped silica cuprous sulphate (CDSCS) as an efficient heterogeneous nano-catalyst, *J. Heterocyclic Chem.*, 2016, 54, 1863.
 43. (a) S. Vidyacharan, A. Murugan, D. S. Sharada, C(sp²)–H Functionalization of 2H-Indazoles at C3-Position via Palladium(II)-Catalyzed Isocyanide Insertion Strategy Leading to Diverse Heterocycles *J. Org. Chem.* 2016, 81, 2837. (b) S. Vidyacharan, A. Sagar, N. C. Chaitra, D. S. Sharada, A facile synthesis of 2H-indazoles under neat conditions and further transformation into aza-γ-carboline alkaloid analogues in a tandem one-pot fashion, *RSC Adv.* 2014, 4, 34232. (c) S. Vidyacharan, C. Adhikari, V. S. Krishna, R. S. Reshma, D. Sriram, D. S. Sharada, A robust synthesis of functionalized

- 2H-indazoles via solid state melt reaction (SSMR) and their anti-tubercular activity. *Bioorg. Med. Chem. Lett.*, 2017, 27, 1593.
44. J. Hu, Y. Cheng, Y. Yang, Y. Rao, A general and efficient approach to 2H-indazoles and 1H-pyrazoles through copper-catalyzed intramolecular N–N bond formation under mild conditions, *Chem. Commun.* 2011, 47, 10133.
 45. M. Akazome, T. Kondo, Y. Watanabe, Palladium Complex-Catalyzed Reductive N-Heterocyclization of Nitroarenes: Novel Synthesis of Indole and 2H-Indazole Derivatives, *J. Org. Chem.* 1994, 59, 3375.
 46. (a) J. I. G. Cadogan, R. K. Mackie, 2-Phenylindazole - *Organic Syntheses, Org. Synth.* 1968, 48, 113. (b) J. Pérez-Villanueva, L. Yépez-Mulia, I. González-Sánchez, J. F. Palacios-Espinosa, O. Soria-Arteche, T. R. Sainz-Espuñes, M. A. Cerbón, K. Rodríguez-Villar, A. K. Rodríguez-Vicente, M. Cortés-Gines, Z. Custodio-Galván, D. B. Estrada-Castro, Synthesis and Biological Evaluation of 2H-Indazole Derivatives: Towards Antimicrobial and Anti-Inflammatory Dual Agents, *Molecules* 2017, 22, 1864.
 47. N. E. Genung, L. Wei, G. E. Aspnes. Regioselective Synthesis of 2H-Indazoles Using a Mild, One-Pot Condensation–Cadogan Reductive Cyclization, *Org. Lett.*, 2014, 16, 3114.
 48. T. V. Nykaza, T. S. Harrison, A. Ghosh, R. A. Putnik, A. T. Radosevich, A Biphilic Phosphetane Catalyzes N–N Bond-Forming Cadogan Heterocyclization via PIII/PV=O Redox Cycling, *J. Am. Chem. Soc.* 2017, 139, 6839.
 49. A. K. Saikia, R. Unnava, K. Indukuri, S. Sarkar. Regioselective one-pot, three-component synthesis of substituted 2H-indazoles from 2-nitroarylaldehyde, alkyne and amine catalyzed by the CuBr/Zn(OTf)₂ system, *RSC Adv.* 2014, 4, 55296.
 50. (a) N. Sudhapriya, A. Nandakumar, P. T. Perumal. Facile synthesis of 2-substituted quinolines and 3-alkynyl-2-aryl-2H-Indazole via SnCl₂-mediated reductive cyclization, *RSC Adv.* 2014, 4, 58476. (b) N. Sudhapriya, C. Balachandran, S. Awale, P. T. Perumal, Sn(II)-Mediated facile approach for the synthesis of 2-aryl-2H-indazole-3-phosphonates and their anticancer activities, *New J. Chem.* 2017, 41, 5582.
 51. (a) G. H. Ahn, J. J. Lee, Y. M. Jun, B. M. Lee, B. H. Kim, Reductive heterocyclizations via indium–iodine-promoted conversion of 2-nitroaryl imines or 2-nitroarenes to 2,3-diaryl-substituted indazoles, *Org. Biomol. Chem.* 2007, 5, 2472. (b) F. Sun, X. Feng, X. Zhao, Z.-B. Huang, D.-Q. Shi, An efficient synthesis of 2H-indazoles via reductive cyclization of 2-nitrobenzylamines induced by low-valent titanium reagent *Tetrahedron*, 2012, 68, 3851. (c) H. B. Abed, J. Schoene, M. Christmann, M. Nazare, *Contemporary Organic Synthesis Org. Biomol. Chem.*, 2016, 14, 8520. (d) T.Y. Wu, S. Dhole, M. Selvaraju, C.M. Sun, Regioselective Synthesis of Pyranone-Fused Indazoles via Reductive Cyclization and Alkyne Insertion, *ACS Comb. Sci.* 2018, 20, 156.
 52. C. Wu, Y. Fang, R. C. Larock, F. Shi, A Facile, One-Pot Synthesis of β -Substituted (Z)-Acrylonitriles Utilizing an α -Diaminoboryl Carbanion, *Org. Lett.*, 2010, 12, 2171.
 53. B. Haag, Z. Peng, P. Knochel, Preparation of Polyfunctional Indazoles and Heteroarylazo Compounds Using Highly Functionalized Zinc Reagents, *Org. Lett.*, 2009, 11, 4270.
 54. J. J. Song, N. K. Yee, A Novel Synthesis of 2-Aryl-2H-indazoles via a Palladium-Catalyzed Intramolecular Amination Reaction, *Org. Lett.*, 2000, 2, 519.
 55. K. Y. Lee, S. Gowrisankar, J. N. Kim, Facile synthesis of 2H-indazole derivatives starting from the Baylis–Hillman adducts of 2-cyclohexen-1-one, *Tetrahedron Lett.*, 2005, 46, 5387.
 56. N. Halland, M. Nazare, O. R'kyek, J. Alonso, M. Urmann, A. Lindenschmidt, A general and mild palladium-catalyzed domino reaction for the synthesis of 2H-indazoles, *Angew. Chem., Int. Ed.* 2009, 48, 6879.
 57. W. Wei, X. Li, M. Gu, H. Yao, A. Lin, Cu/Pd cooperatively catalyzed tandem C–N and C–P bond formation: access to phosphorylated 2H-indazoles, *Org. Biomol. Chem.* 2017, 15, 8458.
 58. a) X. Yi, L. Jiao, C. Xi, *Org. Biomol. Chem.* 2016, 14, 9912. (b) X. Yi, C. Xi, Iodine-catalyzed aerobic oxidation of o-alkylazoarenes to 2H-indazoles, *Tetrahedron* 2017, 73, 1311.
 59. (a) Y. Lian, R. G. Bergman, L. D. Lavis, J. A. Ellman, Rhodium(III)-Catalyzed Indazole Synthesis by C–H Bond Functionalization and Cyclative Capture, *J. Am. Chem. Soc.* 2013, 135, 7122. (b) H. Li, P. Li, L. Wang, Direct Access to Acylated Azobenzenes via Pd-Catalyzed C–H Functionalization and

- Further Transformation into an Indazole Backbone, Org. Lett.* 2013, 15, 620. (c) J. R. Hummel, J. A. Ellman, *J. Am. Chem. Soc.* 2015, 137, 490. (d) X. Geng, C. Wang, *Rhenium-Catalyzed [4 + 1] Annulation of Azobenzenes and Aldehydes via Isolable Cyclic Rhenium(I) Complexes, Org. Lett.* 2015, 17, 2434. (e) T. Jeong, S. H. Han, S. Han, S. Sharma, J. Park, J. S. Lee, J. H. Kwak, Y. H. Jung, I. S. Kim, *Access to 3-Acyl-(2H)-indazoles via Rh(III)-Catalyzed C–H Addition and Cyclization of Azobenzenes with α -Keto Aldehydes, Org. Lett.,* 2016, 18, 232.
60. Z. Long, Z. Wang, D. Zhou, D. Wan, J. You, *Rh(III)-Catalyzed Regio- and Chemoselective [4 + 1]-Annulation of Azoxy Compounds with Diazoesters for the Synthesis of 2H-Indazoles: Roles of the Azoxy Oxygen Atom, Org. Lett.,* 2017, 19, 2777.
61. Z. Long, Y. Yang, J. You, *Rh(III)-Catalyzed [4 + 1]-Annulation of Azoxy Compounds with Alkynes: A Regioselective Approach to 2H-Indazoles, Org. Lett.,* 2017, 19, 2781.
62. (a) B. Avila, M. H. El-Dakdouki, M. Z. Nazer, J. G. Harrison, D. J. Tantillo, M. J. Haddadin, M. J. Kurth, *Acid and base catalyzed Davis–Beirut reaction: experimental and theoretical mechanistic studies and synthesis of novel 3-amino-2H-indazoles, Tetrahedron Lett.* 2012, 53, 6475. (b) J. S. Zhu, M. R. Duong, A. P. Teuthorn, J. Y. Lu, J. H. Son, M. J. Haddadin, M. J. Kurth, *Davis–Beirut Reaction: Alkoxide versus Hydroxide Addition to the Key o-Nitrosoimine Intermediate, Org. Lett.* 2018, 20, 1308.
63. A. Taher, S. Ladwa, S. T. Rajan, G. W. Weaver, *Synthesis of 2-aryl-2H-indazoles by base catalysed reaction of 2-nitrobenzyl triphenylphosphonium bromide and aryl isocyanates, Tetrahedron Lett.* 2000, 41, 9893.
64. B. J. Stokes, C. V. Vogel, L. K. Urnezis, M. Pan, T. G. Driver, *Intramolecular Fe(II)-Catalyzed N–O or N–N Bond Formation from Aryl Azides, Org. Lett.* 2010, 12, 2884.