[Pamwar, 9(6): June 2022]

ISSN 2348 - 8034 Impact Factor- 5.070

# GLOBAL JOURNAL OF ENGINEERING SCIENCE AND RESEARCHES STUDY OF REMEDIABLE SUBSTANCES PRESENT IN KANH RIVER OF INDORE

Minakshi Panwar\*1, Devraj Pranami², and Deeksha Raghuvanshi³

\*1,2&3 Department of Chemical Sciences, Christian Eminent College, Indore (M.P.) India

#### **ABSTRACT**

The city was founded along the banks of the Kanh River by the Holkar dynasty. It was an outcome of the Mandsaur. Peace Treaty signed between the East India Company and Maharaja Malhar Rao Holkar in 1818," informs Kishore Kodwani, a social activist in Indore who is working for the purification of the Kanh. We can find pictures of people bathing in the Kanh water up to the decade of the 50s. Religious rituals were regularly performed alongside the river. The problem began in the 60s and the 70s. It was the time when Indore witnessed a rapid wave of industrialisation<sup>1</sup>, <sup>2,4</sup>. The water obtained from various resources such as underground (Well water, Boar well water), surface water (rivers, ponds, lakes, marine water) and canals of tribal area of Madhya Pradesh, having excess amount of organic and inorganic components, which are hazardous to human being and animals<sup>1</sup>

**Keywords**: metallurgy, purification, electrovalent, synthesis, Interference, metallurgy, purification, hardness, turbidity.

#### I. INTRODUCTION

Few cared for the burden of the waste that this development produced. It eventually went into the rivers, as the infrastructure remained underdeveloped. Like the size of the city, the volume of the sewage disposed into the Kahn river is only increasing with time."In the name of infrastructure, we built the ring road, several bypasses, new colonies, parks and The Bus Rapid Transit System (BRTS) or Ahilya Path. However, the waste management was left to the Kahn river. The very river that was the reason for the foundation of the city is now throttled by its growth<sup>5</sup> One of world's largest religious gatherings-SimhasthaKumbha- is held after every 12 years on the banks of the Kshipra at Ujjain. Reduced to a black nallah for the non-rainy part of the year, the Kahn river is the biggest source of contamination for the shipra. An overwhelming majority of industrial waste is dumped into its 21-kilometre stretch that flows through the urban areas. Saraswati, a tributary that emanates from the same mountain, joins the Kahh at Indore<sup>1,2</sup>.

The data released by the Indore Municipal Corporation estimate the daily volume of sewage from Indore at around 270 million liters per day (MLD). The combined total capacity of the existing sewage treatment plants is not more than 90 MLD. In other words, approximately 190 MLD of untreated sewage is added each day to the Kahn river, which eventually pours into the Shipra. The city of Ujjain gets its water supply from the same polluted river.

## II. MATERIALS AND METHODS

pH: pH can be viewed as an abbreviation for power of hydrogen or more completely , power of the concentration of hydrogen ion. Most natural water is alkaline in nature due to presence of bicarbonates and carbonates formed due to dissolution of atmospheric Carbon dioxide. pH can be drastically change due to prevailing biochemical activities undergoing in water. Photosynthetic activity increases the pH due to consumption of free CO2 and dissociation of bicarbonates into carbonates. The carbonate are much stronger alkalise than the bicarbonates H: pH can be viewed as an abbreviation for power of hydrogen or more completely , power of the concentration of hydrogen ion. Most natural water is alkaline in nature due to presence of bicarbonates and carbonates formed due to dissolution of atmospheric Carbon dioxide. pH can be drastically change due to prevailing biochemical activities undergoing in water. Photosynthetic activity increases the pH due to consumption of free CO2 and dissociation of bicarbonates into carbonates. The carbonate are much stronger alkalise than the bicarbonates 4 H: pH can be viewed as an abbreviation for power of hydrogen or more completely , power of the concentration of hydrogen ion. Most natural water is alkaline in nature due to presence of bicarbonates and carbonates formed due to dissolution of atmospheric





## [Pamwar, 9(6): June 2022]

ISSN 2348 - 8034 Impact Factor- 5.070

Carbon dioxide. pH can be drastically change due to prevailing biochemical activities undergoing in water. Photosynthetic activity increases the pH due to consumption of free CO2 and dissociation of bicarbonates into carbonates. The carbonate are much stronger alkalise than the bicarbonates 4

The sewage is not the only problem afflicting the Kahn river. The mindless encroachment of its catchment area and 33 slum colonies have swallowed a great part of the river. Its flow has greatly reduced because of the illegal constructions cropping in Kahn's catchment area. The human waste produced from the slum colonies also flows into the Kahn river. Besides, illegal colonies have mushroomed along the banks of the river. As a result, the river has changed course at some locations. People have erected retaining walls on others, further restricting its natural flow<sup>6</sup>.

# Table for experimental parameters

Parameters	Values obtained
pH	6.63 without turbidity
<b>pH</b> - Neutral water has a pH of 7, acidic solutions	6.73 with turbidity
have values between 0-6 and alkaline solutions have	
values between 8-14	
Alkalinity	Between 80 to 100 mg/L as CaCO <sub>3.</sub>
<b>Alkalinity</b> is a measure of the buffering capacity of	
water, or the capacity of the water to neutralise acids	
and resist pH change.	
<b>Turbidity</b> is a measure of the cloudiness or haziness	Highly turbid
in water caused by suspended solids (eg sediment,	
algae). Turbidity is expressed in Nephelometric	
Turbidity Units (NTU) and is measured using a	
relationship of light reflected from a given sample.	
Hardness is a soap consuming capacity of water	Total Hardness- 410ppm
hence if water is hard it surly contain any or more	Calcium Hardness- 326 ppm
substance out of the following ;- CaCO <sub>3</sub> , CaCO <sub>3</sub> ,	
CaCl <sub>2</sub> , MgCl <sub>2</sub> , CaSO <sub>4</sub> , MgSO <sub>4</sub> ,CaHCO <sub>3</sub> ,	
MgHCO <sub>3</sub> .	

## III. RESULT AND DISCUSSION

The characteristics which render the stream water unacceptable to physical senses are included in this group. Some of the common and important characteristics are enumerated here under:

## pН

The value indicates the ionic equilibrium. It indicates the acidic or alkaline nature of water. A few industries discharge acidic effluents while other discharge alkaline. Natural water body has a tendency to resist the input of acid and alkali. This is known as buffering tendency. This does not allow change in the pH of the water. If it does not so then change in the pH affects the life of the biotic molecules and the accessibility of the few ions.

#### **Turbidity**

The crystal clear water is always consider as good quality water. Clarity of water is an important and essential characteristic in producing the product design for human consumption and in many manufacturing uses. The clarity of natural body of water is major determinant of the condition.

## **Suspended Solids**

Presence of heavy swarm of suspended and floating material in water bodies makes their look highly rough and untidy. The changes brought in chemical characteristics through the domestic and industrial effluent cause pollution. It can thus be used as parameter to indicate the pollution.





[Pamwar, 9(6): June 2022]

ISSN 2348 - 8034 Impact Factor- 5.070

#### **Hardness**

The hardness of water is its soap consuming capacity. Hardness is a measure of an aggregate property of water. Hardness of many surfaces water is primarily a function of carbonate, bicarbonate and hydroxide content, it is taken as indication of concentration of these constituents.

#### **Survey of Literature**

Fluoride and other Elements

The features of this organic fraction, which occupies the size range generally defined as colloidal (> 1 nm to < 0.4  $\mu$ m), are the ability to complex or sorbs other chemical species from solution, the ability to serve as reducing reagents in chemical reactions, and the ability to either enhance or retard photochemical reactions occurring in solution. Out of Hazardous component Fluoride is one of them. Fluoride has interfering property i.e. it interferes in any type of analysis. Interfering radicals are tartrate, fluoride, borate, phosphate, oxalate; they are also known as anionic radicals. They are usually used to form a complex with 3rd group elements. Aluminium, Chromium and Iron are the basic radicals of this third group and it is well known fact that radical fluorination is a type of fluorination reaction, complementary to nucleophilic and electrophonic approaches. Fluoride has an inherent property to interfere in any type of analysis. Most of the water bodies viz. rivers, ponds canals of tribal area of Madhya Pradesh, having excess amount of fluoride. Previous studies and researches show that this excess fluoride affected the human health by causing several diseases<sup>8</sup>.

### IV. CONCLUSION

All facts of this paper reveal that for the researchers, there is a new criteria focusing on the remediable organic and inorganic components present in kanh and other tributary river water and it's important to remove interference of excess of these substances obtained during enormous industrial processes such as manufacturing, purification, synthesis and metallurgy etc. and all the component whether the organic or inorganic in nature can spoil the drinking and other consumption quality of water hence that will cause ill effect on health of living beings, that is why prior to treat or remove. The results obtained from the observations will be of interest of analysts and researchers as well.

### **REFERENCES**

- 1. C. Sigleo ,J. C. Means, Reviews of Environmental Contamination and Toxicology pp 123-147.
- 2. Abelson PH, Hare PE (1970) Uptake of amino acids by kerogen. Carnegie Inst Wash Yearb 69:297–303. Google Scholar
- 3. Abelson PH, Hare PE (1971) Reactions of amino acids with natural and artificial humus and kerogens. Carnegie Inst Wash Yearb 70:327–333. Google Scholar
- 4. Boehm PD, Quinn JG (1973) Solubilization of hydrocarbons by the dissolved organic matter in sea water. GeochimCosmochimActa 37:2459–2477.CrossRefGoogle Scholar
- 5. Boynton WR, Kemp WM, Keefe CW (1982) A comparative analysis of nutrients and other factor influencing estuarine phytoplankton production. In: Kennedy VC (ed) Estuarine Comparisons. Academic Press, NY, pp 69–90.Google Scholar
- 6. Home » India news » The story of Kahn: A river swallowed by the spreading city of Indore
- 7. Brownawell BJ (1986) The role of colloidal organic matter in the marine geochemistry of PCBs. PhD Thesis Massachusetts Institute of Technology/Woods Hole Oceanographic Institution, MA. p 318.
- 8. E. Lombi, R.E. Hamon, in Encyclopedia of Soils in the Environment, 2005.

